

432-1361

12/11/2013

11/5

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Bayer Environmental Science
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

DEC 11 2013

Subject: Amended label adding pollinator protection language
Product Name: Merit 60 WSP Greenhouse and Nursery Insecticide
EPA Reg. No. 432-1361
EPA Decision No. 485406
Submission dated September 27, 2013

Dear Dr. Pangilinan:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable. A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. See 40 CFR 156.10(a)(6).

Under 40 CFR 152.130(d), EPA may establish dates by which all product distributed or sold by the registrant must bear revised labeling. The following paragraphs set forth the schedule for ensuring that that your product bears revised labeling within a reasonable time period.

- Any product released for shipment after 2/28/14 must bear the new label.

If these conditions are not complied with, EPA will take appropriate action against this registration. If you have any questions please contact Dr. Jennifer Urbanski at 703-347-0156 or urbanski.jennifer@epa.gov.

Regards,
A handwritten signature in black ink, appearing to read "Venus Eagle", followed by a long horizontal line.
Venus Eagle, Product Manager (01)
Insecticide-Rodenticide Branch
Registration Division (7505P)

2/15

GROUP

4A

INSECTICIDE

Merit[®] 60 WSP Greenhouse and Nursery Insecticide

For foliar and systemic insect control on ornamental and vegetable transplants in greenhouses and nurseries.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine 60.0%

OTHER INGREDIENTS: 40.0%

TOTAL: 100.0%

Keep water soluble packets in this container and store in a cool dry place but not below freezing (32F).

Do Not Remove Packets From Container Except For Immediate Use

EPA Reg. No. 432-1361

EPA Est. No. ____

STOP - READ THE LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCION AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
In case of emergency call toll free the Bayer Environmental Science Emergency Response Telephone No. 1-800-334-7577. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.	
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	

ACCEPTED

DEC 11 2013

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

EPA. Reg. No: 432-1361

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS:

- When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.


ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on the foliage of blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT
BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS
FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



Look for the bee hazard icon in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

SEE INDIVIDUAL CROPS FOR SPECIFIC POLLINATOR PROTECTION APPLICATION RESTRICTIONS. IF NONE EXIST UNDER THE SPECIFIC CROP, FOR OUTDOOR FOLIAR APPLICATIONS, FOLLOW THESE APPLICATION DIRECTIONS FOR FOOD/FEED CROPS AND COMMERCIALY GROWN ORNAMENTALS THAT ARE ATTRACTIVE TO POLLINATORS.

FOR FOOD/FEED CROPS AND COMMERCIALY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS



Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
- The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify beekeepers no less than 48-hours prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

PRODUCT INFORMATION

Merit® 60 WSP Greenhouse and Nursery Insecticide is for insect control on ornamental and vegetable plants intended for resale (transplants) in nurseries and greenhouses. Merit 60 WSP Greenhouse and Nursery Insecticide may be applied either as a spray or as a drench. Thorough coverage is required when applied as a spray. Merit 60 WSP Greenhouse and Nursery Insecticide is a systemic product and will be translocated upward within the plant when applied as a drench. To assure optimum effectiveness, the product must be placed where the roots of the plant can absorb the active ingredient. Irrigate moderately but thoroughly after application, allowing no leaching and runoff from container for at least three irrigations or 10 days whichever ever is longer.

Resistance: Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, conform the use of this product to resistance management strategies established for the use area.

Merit 60 WSP Greenhouse and Nursery Insecticide contains a Group 4A insecticide. Insect biotypes with acquired or inherent resistance to Group 4A insecticides may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

The active ingredient in Merit 60 WSP Greenhouse and Nursery Insecticide is a member of the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to Merit 60 WSP Greenhouse and Nursery Insecticide.

Contact your Cooperative extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

For resistance management purposes, do not make a foliar application with any chloronicotinyl insecticide following a Merit 60 WSP Greenhouse and Nursery Insecticide soil application in the same crop.

Incorporation: Incorporation of Merit 60 WSP Greenhouse and Nursery Insecticide can be achieved by cultivation, irrigation, rainfall, mechanical placement, soil injection, drenching, and broadcast sprays.

Woody Perennials: Onset of protection is slower than in herbaceous species. A delay of 2 or more weeks will be expected. Longer delays may be expected with larger plants. Therefore, make application well in advance of expected insect activity.

Bark Media: Media with 30 to 50% or more bark content may confer a shorter period of protection when treated with Merit 60 WSP Greenhouse and Nursery Insecticide.

Tank Mixes: Merit 60 WSP Greenhouse and Nursery Insecticide has been found to be compatible with commonly used liquid fertilizers, fungicides and insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

RESTRICTION: Do not use Merit 60 WSP Greenhouse and Nursery Insecticide packets in a tank mix with products that contain boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water. However, normal chlorinated water is acceptable for mixing.

MIXING: Within each foil pouch is a clear inner packet containing Merit 60 WSP Greenhouse and Nursery Insecticide. The clear inner packet is water soluble. Do not allow packets to become wet prior to adding to the tank. Do not handle the clear inner packets with wet hands or wet gloves. Rough handling may cause breakage. Reseal outer carton to protect remaining packets.

To prepare the mixture, remove the outer foil pouch and drop the required number of unopened clear water soluble packets into the spray tank while filling the tank with the desired amount of water. The required number of packets is specified under 'Dose' in the tables below. Operate the agitator while mixing. Depending on the water temperature and the degree of agitation, the packets will completely dissolve within a few minutes from the time they are added to the water.

APPLICATION TO GRASSY AREAS IN NURSERIES

Merit 60 WSP Greenhouse and Nursery Insecticide can be used for the control of soil inhabiting pests of grassy areas of nurseries, such as northern and southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; European chafer, *Rhizotrogus majalis*; green June beetle, *Cotinis nitida*; May or June beetle, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; Oriental beetle, *Anomala orientalis*; sugarcane beetle, *Euethola rugiceps humilis*; billbugs, *Sphenophorus* spp.; annual bluegrass weevil, *Listronotus maculicollis*; black turfgrass ataenius, *Ataenius spretulus*; *Aphodius* spp. crane flies, *Tipula* spp.; and mole crickets, *Scapteriscus* spp. Merit 60 WSP Greenhouse and Nursery Insecticide can also be used for suppression of cutworms and chinchbugs. Merit 60 WSP Greenhouse and Nursery Insecticide can be used as directed on nursery grass in sites such as under or around field or container grown plants, on roadways or other grassy areas in or around nurseries. Merit 60 WSP Greenhouse and Nursery Insecticide cannot be used on commercial sod farms.

The active ingredient in Merit 60 WSP Greenhouse and Nursery Insecticide has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Do not apply to grassy areas if the soil is waterlogged or saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Application cannot exceed a total of 10.7 oz (0.4 lb of active ingredient) per acre per year.

Application Equipment for Use on Grassy Areas in Nurseries: Apply Merit 60 WSP Greenhouse and Nursery Insecticide in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for soil

application of insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.

APPLICATION THROUGH IRRIGATION SYSTEMS

Apply Merit 60 WSP Greenhouse and Nursery Insecticide at rates indicated on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:10 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply Merit 60 WSP Greenhouse and Nursery Insecticide only through microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from nonuniform distribution of treated water.

If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments when the need arises.

SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

DIRECTIONS FOR FOLIAR APPLICATION TO GRASSY AREAS OF NURSERIES		
SITE	PEST	DOSE
Grassy areas of Field & Forest Nurseries	Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataenius Cutworms (suppression) European chafer	Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer
	Chinch bugs (suppression)	Mole crickets
		One packet per 2,900-3,850 sq ft
		One packet per 2,900 sq ft
<p>For optimum control of grubs, billbugs and annual bluegrass-weevil, make application prior to egg hatch of the target pest. Be sure to read "APPLICATION EQUIPMENT" Section of this label.</p> <p>For suppression of chinch bugs, make application prior to the hatching of the first instar nymphs.</p> <p>For optimum control of mole crickets, make application prior to or during the period of peak egg hatch. If application is not made prior to egg hatch and adults or large nymphs are present and actively tunneling, then apply Merit 60 WSP Greenhouse and Nursery Insecticide in combination with a remedial insecticide to minimize damage to turf. Follow label instructions for other insecticides when tank-mixing.</p> <p>Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application. For optimal control, if rainfall does not occur within 24 hours after application, irrigate to move the active ingredient through the thatch. Mow grass area only after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.</p>		
RESTRICTIONS: <ul style="list-style-type: none"> Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per year. Do not allow this product to contact plants in bloom while bees are foraging the treatment area. Do not graze treated areas or use clippings for treated areas for feed or forage. Avoid runoff or puddling of irrigation water following application. Do not apply Merit 60 WSP Greenhouse and Nursery Insecticide to soils which are water logged or saturated, which will not allow penetration into the root zone of the plants. 		

DIRECTIONS FOR APPLICATIONS TO ORNAMENTALS AND VEGETABLE TRANSPLANTS		
For foliar and systemic insect control in and around field grown nurseries, outdoor containerized ornamentals, ornamentals grown in greenhouses and ornamentals grown in flats, benches or beds, for vegetable transplants and resale.		
CROP	PEST	DOSE
Trees (Non-bearing fruit and Nut trees*) Shrubs Evergreens Flowers Foliage plants Ground covers Vegetable transplants**	Adelgids Aphids Japanese beetles (adults) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter)	Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies
	One packet per 100 gal water	
	Foliar Applications: Start treatments prior to establishment of high pest populations and reapply on an as needed basis. For resistance management purposes, do not make a foliar application with Merit 60 WSP Greenhouse and Nursery Insecticide following a Merit 60 WSP Greenhouse and Nursery Insecticide soil application in the same crop.	
	White grub larvae (such as Japanese beetle larvae, chafers, <i>Phyllophaga</i> spp. Asiatic garden beetle, Oriental beetle)	One packet per 2,900-3,850 sq ft.
	Broadcast Applications: Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1,000 sq. ft. Apply specified dosage in enough water to ensure Merit 60 WSP Greenhouse and Nursery Insecticide reaches the root zone but does not leach out of the container. Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is irrigated thoroughly after application. * Non-bearing fruit and nut trees are those trees that will not bear fruit or nuts for one year after application. ** Only for use on the following vegetable plants intended for resale including: Broccoli, Chinese broccoli, Broccoli raab, Brussels sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground cherry, Kale, Kohlrabi, Lettuce, Mustard greens, Pepinos, Peppers, Potatoes, Rape greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.	
RESTRICTIONS FOR OUTDOOR USE:		
<ul style="list-style-type: none">Follow application restrictions FOR COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS indicated on page [x] to protect bees and other insect pollinators.Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per year.Do not apply this product, by any application method, to linden, basswood, or other Tilia species.		

DIRECTIONS FOR DRENCH AND IRRIGATION APPLICATIONS

For use only in greenhouses and on nursery ornamentals and vegetable transplants using soil drenches, micro-irrigation, drip irrigation, overhead irrigation, or hand-held or motorized calibrated irrigation equipment.

PEST	USE PATTERN		DOSE - MERIT 60 WSP		APPLICATION INSTRUCTIONS
Adelgids	Plants in containers	Herbaceous species - including vegetables and transplants ⁵ (one or two plants per pot)	Container size (inches)	No. pots treated with one (1) Packet	Evenly distribute one (1) packet of Merit 60 WSP in the stated number of pots, using sufficient water volume to wet potting medium without loss of liquid through leaching. Apply according to label directions. Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient due to leaching.
Aphids				3,000	
Armored scale (suppression)			2	2,000	
Fungus gnat (larvae only) ¹			3	1,500	
Japanese beetles (adults)			4	1,200	
Lacebugs			5	1,000	
Leaf beetles			6	850	
(including elm and viburnum leaf beetles)			7	750	
Leafhoppers (including glassy-winged sharpshooter)			8	675	
Leafminers			9	600	
Mealybugs			10	550	
Psyllids			11	500	
Root mealybugs ²			12	450	
Root weevil complex (such as Black vine weevil, Apopka weevil, Citrus root weevil ³)	Woody perennial species	2	2,000		
Soft scale		3	1,350		
Thrips (suppression) ⁴		4	1,000		
White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)		5	800		
Whiteflies		6	650		
		7	550		
		8	500		
		9	450		
		10	400		
		11	350		
		12	300		
		Herbaceous species - including vegetable transplants ⁵ (three or more plants per pot)	Use the above woody perennial species rates		
	Ornamental and vegetable ⁵ plants grown in flats, benches, or beds	One packet per 3,000 sq ft		Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of mixture per 1,000 sq ft. Apply as a broadcast treatment and incorporate into the medium before planting or apply after plants are established. If application is made to established plants, optimum control will be attained if area is lightly irrigated after application. Allow no leaching or runoff for 10 days after application.	
	Containerized plants	Container size	No. pots treated with one (1) packet	Apply in sufficient water to wet the potting medium. For optimum control, make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.	
		1 gallon	240 to 120		
		2 gallon	120 to 60		
		3 gallon	90 to 40		
		5 gallon	65 to 30		

RESTRICTIONS FOR OUTDOOR USES:

- Follow application restrictions FOR COMMERCIAL GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS indicated on page [x] to protect bees and other insect pollinators.
- Do not apply to soils which are water-logged or saturated, which will not allow the penetration of the insecticide into the root zone of the plants.
- Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per year.
- Do not apply this product, by any application method, to linden, basswood, or other Tilia species.

- ¹ **Fungus gnat larvae** in the soil will be controlled by drench or incorporation. **No adult fungus gnat control.** Other foliar insect control is achieved by the uptake of Merit 60 WSP Greenhouse and Nursery Insecticide from a healthy root system translocating the active ingredient up into the plant.
- ² **Root mealybug** control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: one (1) packet in 150 gallons of water.
- ³ **Citrus root weevil:** For use on non-bearing citrus nursery stock.
- ⁴ **Thrips** suppression on foliage only. Thrips in buds and flowers will not be suppressed.
- ⁵ For use only on the following vegetable plants (intended for transplant resale): Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

DIRECTIONS FOR SOIL APPLICATIONS TO FIELD GROWN NURSERY STOCK			
PEST	USE PATTERN	DOSAGE	INSTRUCTION
White grub larvae (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	Field and forest nursery stock	Apply as a uniform band to the ground on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap. Use one (1) packet per 1,000 ft of row or 3,000 sq ft.	Apply May through July. For optimum control, treatment must be followed by rainfall or irrigation. Do not use less than 2 gallons of spray volume per 1,000 square feet.
RESTRICTIONS: <ul style="list-style-type: none"> Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per year. Do not allow this product to contact plants in bloom while bees are foraging the treatment area. Do not apply Merit 60 WSP Greenhouse and Nursery Insecticide to soils which are water logged or saturated, which will not allow penetration into the root zone of the plants. 			

DIRECTIONS FOR SOIL DRENCH AND SOIL INJECTION FOR TREES AND SHRUBS FOR NURSERIES AND GREENHOUSES				
Adelgids	Eucalyptus longhorned borers	Leaf beetles (including elm and viburnum leaf beetles)	Pine tip moth larvae	Soft scales
Aphids	Flatheaded borers (including bronze birch, emerald ash, and alder borers)	Leafhoppers (including glassy-winged sharpshooter)	Psyllids	Thrips (suppression)
Armored scales (suppression)	Japanese beetles (adults)	Leafminers	Royal palm bugs	White grub larvae
Black vine weevil larvae	Lacebugs	Mealybugs	Sawfly larvae	Whiteflies
Trees	1 packet (20 grams) per 8 to 16 inches of cumulative trunk diameter			
Soil Injection: GRID SYSTEM: Space holes on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.				
Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. For optimum control, keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.				
No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.				
Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.				
For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.				
Shrubs	1 packet (20 grams) per 8 to 16 feet of cumulative shrub height			
Soil Injection: Apply to individual plants using dosage indicated.				
Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per shrub.				
No Soil Injection Application Allowed in Nassau or Suffolk Counties of New York.				
Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.				
GROUND TREATMENT PRE-PLANTING FOR FLOWERS AND GROUNDCOVERS	1 packet (20 grams) per 3,000 sq ft			
Apply as a broadcast treatment and incorporate into the soil before planting.				
Restrictions for Outdoor Uses:				
<ul style="list-style-type: none">Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per yearDo not apply to soils which are water-logged or saturated, which will not allow the penetration of the insecticide into the root zone of the plants.For use only on non-bearing fruit and nut trees. Non-bearing fruit and nut trees are those trees that will not bear fruit or nuts for one year after application.Do not apply this product, by any application method, to linden, basswood, or other Tilia species.				

MERIT 60 WSP - Ebb & Flood Application

Merit 60 WSP Greenhouse and Nursery Insecticide may be applied through Ebb and Flood applications. To assure accurate uptake prior to treatment, bring a minimum of 10 plants up to a known field capacity and allowed to dry out for one or two days. Re-wet these plant to determine how much water on average each plant will absorb to bring it back at field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum to flood your smallest treatment area. Do not allow leachate run out for first 10 days after application, in order to retain the product and facilitate full plant uptake of the active ingredient. This will minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

DIRECTIONS FOR EBB AND FLOOD APPLICATIONS IN GREENHOUSES AND NURSERIES

Adelgids	Japanese beetles	Leafminers	Root mealybugs ²	Thrips
Aphids	(adults)	Mealybugs	Root weevil complex:	(suppression) ⁴
Armored scales	Lacebugs	Psyllids	(such as Apopka	Whiteflies
(suppression)	Leaf beetles		weevil, Black vine	White grub larvae (such
Fungus gnats	(including elm and		weevil, Citrus Root	as Japanese beetle,
(larvae only) ¹	viburnum leaf beetles)		weevil ³)	masked chafers,
	Leafhoppers (including		Soft scales	European chafer,
	glassy-winged			Oriental beetle, Asiatic
	sharpshooter)			garden beetle)

Pot sizes (inches)	Number of pots treated with one packet	
	Herbaceous species including vegetable-transplants ⁵ (one or two plants per pot)	Woody perennials, herbaceous species including vegetable transplants ⁵ (3 or more per pot)
2	3,000	2,000
3	2,000	1,350
4	1,500	1,000
5	1,200	800
6	1,000	650
7	850	550
8	750	500
9	675	450
10	600	400
11	550	350
12	500	300

¹ Fungus gnat larvae in the soil will be controlled by drench or incorporation. No adult Fungus gnat control. Other foliar insect control is achieved by the uptake of Merit 60 WSP Greenhouse and Nursery Insecticide from a healthy root system translocating the active ingredient up into the plant.

² Root mealybug control will require a thorough drenching of containerized media. Coverage is essential for control while minimizing the amount of leachate. Rate: 1.7 fl oz (50 mL) in 150 gallons of water.

³ Citrus root weevil: For use on non-bearing citrus nursery stock.

⁴ Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.

⁵ For use only on the following vegetable transplants(intended for transplant resale): Broccoli, Chinese Broccoli, Broccoli Raab, Brussels Sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground Cherry, Kale, Kohlrabi, Lettuce, Mustard Greens, Pepinos, Peppers, Potatoes, Rape Greens, Sorghum, Sugarbeets, Tomatillo, and Tomato.

RESTRICTIONS FOR OUTDOOR USE:

- Follow application restrictions FOR COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS indicated on page [x] to protect bees and other insect pollinators.

- Do not apply more than 10.7 oz (0.4 lb of active ingredient) per acre per year

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. After removal of all PVA packets, dispose of empty container in a sanitary landfill, by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides above. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577 or contact Chemtrec at 1-800-424-9300.

15/15

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Bayer CropScience LP's control it is impossible for Bayer CropScience LP to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Net Contents: 4, 20 g (0.7 oz) packets/pouch (2.8 oz/pouch), 6, 50 g (0.7 oz) packets/pouch (120 g or 4.2 oz/pouch)

PRODUCED FOR



Bayer Environmental Science

A DIVISION OF BAYER CROPSCIENCE LP
2 T. W. ALEXANDER DRIVE
RESEARCH TRIANGLE PARK, NC 27709

Merit® 60 WSP Greenhouse and Nursery Insecticide (PENDING) 08-23-2013, 09-26-2013, 09-30-2013, 12-02-2013